


## Definite Integrals, continued....

when  $f$  is non-negative  $a \leq x \leq b$ ,

$\int_a^b f(x) dx$  gives the area under the curve from  $x=a$  to  $x=b$



$$\int_{-1}^2 (2x+2) dx = 9$$

$$\int_{-3}^{-1} (2x+2) dx = -4$$

$$\begin{aligned}\int_{-3}^2 (2x+2) dx &= \int_{-3}^{-1} (2x+2) dx + \int_{-1}^2 (2x+2) dx \\ &= -4 + 9 = 5\end{aligned}$$

$$\int_a^b f(x)dx = \int_a^c f(x)dx + \int_c^b f(x)dx$$

### Properties of Definite Integrals

1.  $\int_a^b Kf(x)dx = K \int_a^b f(x)dx$

2.  $\int_a^b (f(x) \pm g(x))dx = \int_a^b f(x)dx \pm \int_a^b g(x)dx$

3.  $\int_a^a f(x)dx = 0$

4.  $\int_a^b f(x)dx = - \int_b^a f(x)dx$

5.

Ex  $\int_0^2 f(x)dx = 4, \int_2^5 f(x)dx = 12, \int_0^2 g(x)dx = -3, \int_0^4 g(x)dx = 6$

a)  $\int_0^2 (3f(x) - g(x))dx = 3(4) - (-3) = 15$

b)  $\int_2^3 g(x)dx + \int_5^6 f(x)dx = 0 - 12 = -12$

c)  $\int_2^4 g(x)dx = \int_0^4 g(x)dx - \int_0^2 g(x)dx = 6 - (-3) = 9$

d)  $\int_{-3}^{-1} \frac{1}{2} f(x+3)dx = \frac{1}{2} \int_{-3}^{-1} f(x+3)dx$   
 $= \frac{1}{2} \int_0^2 f(x)dx = \frac{1}{2}(4) = 2$

Hw 9H p.308 #2-10 even, 11, 12

## 9.4 Fundamental Theorem of Calculus (FTC part 2)

$$\int_a^b f(x) dx = \left[ F(x) \right]_a^b = F(b) - F(a)$$

↑  
Shmoop  
integral  
evaluated on  
the interval  
 $a \leq x \leq b$

↑  
Integral  
value  
at  
 $x=b$

↑  
Integral  
value  
at  
 $x=a$

use a GDC to evaluate

$$\int_0^2 (x^2 + 1) dx \approx 4.67$$

use FTC to evaluate

$$\begin{aligned} \int_0^2 (x^2 + 1) dx &= \frac{1}{3}x^3 + x \Big|_0^2 = \left( \frac{1}{3}(2)^3 + (2) \right) - \left( \frac{1}{3}(0)^3 + (0) \right) \\ &= \frac{8}{3} + 2 \approx 4.67 \end{aligned}$$

Ex Evaluate without GDC

$$\begin{aligned} \text{a) } \int_{-2}^1 (u-1) du &= \left[ \frac{1}{2}u^2 - u \right]_{-2}^1 \\ &= \left[ \frac{1}{2}(1)^2 - 1 \right] - \left[ \frac{1}{2}(-2)^2 - (-2) \right] \\ &= -\frac{1}{2} - 4 = -\frac{9}{2} \end{aligned}$$

$$\text{b) } \int_2^3 \frac{1}{t} dt = \left[ \ln t \right]_2^3 = \ln(3) - \ln(2) = \ln\left(\frac{3}{2}\right)$$

$$\begin{array}{l} u=x^2 \\ du=2x dx \end{array}$$

$$\begin{aligned} \text{c) } \int_1^3 4x^2(x-1) dx &= \int_1^3 (4x^3 - 4x^2) dx \\ &= 4 \left[ \int_1^3 x^3 dx - \int_1^3 x^2 dx \right] \\ &= 4 \left[ \frac{1}{4}x^4 - \frac{1}{3}x^3 \right]_1^3 = \left[ x^4 - \frac{4}{3}x^3 \right]_1^3 \\ &= \left[ 3^4 - \frac{4}{3}(3)^3 \right] - \left[ (1)^4 - \frac{4}{3}(1)^3 \right] = \frac{136}{3} \end{aligned}$$

FTC with more complicated integrals:

$$\begin{aligned} \textcircled{1} \quad \int_1^5 \left[ e^{2x} + \frac{1}{x^2} \right] dx &= \int_1^5 (e^{2x} + x^{-2}) dx \\ &= \left[ \frac{1}{2} e^{2x} - x^{-1} \right]_1^5 = \left( \frac{1}{2} e^{2(5)} - \frac{1}{5} \right) - \left( \frac{1}{2} e^{2(1)} - \frac{1}{1} \right) \\ &= \frac{1}{2} e^{10} - \frac{1}{5} - \frac{1}{2} e^2 + 1 \\ &= \frac{1}{2} e^{10} - \frac{1}{2} e^2 + \frac{4}{5} \end{aligned}$$

$$\begin{aligned}
 ② \quad & \int_0^3 \sqrt{3x+16} dx = \int_0^3 (3x+16)^{1/2} dx \\
 &= \left[ \frac{1}{3} \left[ \frac{2}{3} (3x+16)^{\frac{3}{2}} \right] \right]_0^3 = \left[ \frac{2}{9} (3x+16)^{\frac{3}{2}} \right] \\
 &= \frac{2}{9} (3(3)+16)^{\frac{3}{2}} - \frac{2}{9} (3(0)+16)^{\frac{3}{2}} \\
 &= \frac{2}{9} (25)^{\frac{3}{2}} - \frac{2}{9} (16)^{\frac{3}{2}} \\
 &= \frac{2}{9} \left( (25)^{\frac{3}{2}} - (16)^{\frac{3}{2}} \right)
 \end{aligned}$$

$$\textcircled{3} \quad \int_0^1 (2x^2 + 1)^3 (4x) dx$$

$u = 2x^2 + 1$   
 $du = 4x dx$

$$\int_0^1 u^3 du = \left[ \frac{1}{4} u^4 \right]_0^1$$

HW 9 I P. 310 #1-6, 10  
 HW 9 J 1-4

$$= \frac{1}{4} ( ) - \frac{1}{4} (0)^4$$

$\frac{1}{4} ( ) - \frac{1}{4} (0)^4$   
 mrs. Ruff will undo her  
 damage on Monday